REFERENCES:

[1] IUPAC, **(2009)**, IUPAC Compendium of Chemical Technology- *the Gold Book Heterocyclic compounds,***2.3.2**, 1340.

[2] Prado Silvia R. T.,Cechiel – FilhoV., Buzi F. C., Rogerio C., Silvia Maria Correa S. C., Maria B. M. de Oliveira,**(2004)**, Biological Evaluation of Some Selected Cyclic Imides: Mitochondrical Effects and in vitro Cytotoxicity, *Verlag dev Zeitschbrift fur Nature forshung, Tubingen*, **59c**, 663-672.

[3] Cunha S., RodovolhoW., AzevedoN. R., Marcos de O. M., LaviucciC., VencatoI., **(2002)**, The Michael Reaction of Enaminones with N-(p-tolyl)-maleimide : Synthesis and Structural Analysis of Succinimide - enaminones, *J. Braz. Chem Soc.*,**13(5)**, 629-634.

[4] Papovic-DjordjevicJ. B., DoserMicovicL. I., JuranicI. O., DrakulicB. J.,**(2002)**,Antiproliferative activity of NCI-DTP Glutarimide Derivatives. An alignment independent 3D QSAR study, *J. Serb. Che. Soc.*,**75**, 1167-1179.

[5] Marulasiddaiah R., KalkhambkarR. G., KulkarniM. V., **(2012)**, Synthesis and biological Evaluation of Cyclic Imides with Coumarines Azacoumarines,*Open Journal of Medicinal Chemistry*,**2**, 89-97.

[6] Kale M., PatwardhanK.,**(2013)**,Synthesis of heterocyclic scaffolds with anti-hyperlipiodemic potential: A Review,*Der pharma chemical*,**5(5)**, 213-222.

[7] Journal of Kerbala University, **(2010)**, Synthesis and Evaluation of Biological activity of new cyclic imides derived from 4-aminophenazone 8**(1),** 1-16.

[8] Klotz A. V. and Higgins B. M., **(1991)**, Deamidation and Succinimide Formation by γ N-Methylasparagine: Potential Pitfallsof Amino Acid Analysis, *Archives ofBiochemistry and Biophysics,***291(1)**, 113-120.

[9] Schwendener R.A., Triub T., Schott H., Langhals H., Barth R.F., Groscurth p.,Hengartner H. , **(1989),** Comparative studies of the preparation of immunoliposomeswith the use of two bifunctional coupling agents and investigationof in vitro immunoliposome-target cell binding by cytofluorometryand electron microscopy**,** *Bwchlrmca et Blophystca Actca,* **1026 (1990)**, 69-79

[10] LozanovV., PetrovS., MitevV., **(2003)**, Simultaneous analysis of amino acid and biogenic polyamines byhigh-performance liquid chromatography after pre-column derivatizationwith N-(9-fluorenylmethoxycarbonyloxy)succinimide, *Journal of Chromatography A*, **1025 (2004)** , 201–208

[11] Al-Azzawi A. M., Hamd A. S., **(2011)**,Synthesis, Characterization and Evaluation ofBiological Activity ofNovel Cyclic Imides Containing Heterocycles Based on 2, 5-disubstituted-1,3,4-thiadiazoles, *Al-Anbar J. Vet. Sci.*, **4(2)**, 1-13

[12] TaherpourA. A., MansuriH. R., **(2005)**, Fast Oxidation of Lactams toCyclic Imides UsingMicrowave Irradiation**,** *Turk J Chem*, **29 (2005)**, 317-320.

[13] Allen S. E., SkoogF.,**(1950)**, Stimulation of seedling growth by seed treatments with N-phenyl succinimide derivatives, *Plant Physiology*, 1-5.

[14] Rosana N. S. P., Guilhon Giselle M. S. P., ZoghbiMaria das G. B. ,AraujoI. S., Uetanabaro Ana Paula T. , Santos Lourivaldo S., Brasil Davi do S. B.,**(2013)**, Volatiles, A Glutarimide Alkaloid and Antimicrobial Effects ofCroton pullei (Euphorbiaceae),*Molecules*, **18**, 3195-3205

[15] ZanattaN., Fabio M. da Silva, Andreia M. P. W. da Silva, Aquino E. da C., BonacorsoH. G.,Martins M. A. P., **(2012**), An efficient synthesis of 3-ethoxypyrrolidine-2,5-diones and cis-2,3,3a,6a-tetrahydrofuro[2,3-c]pyrrole-4,6(5H)-diones from *β*-cyanocarboxylic acids, *ARKIVOC*, **(ix)**, 1-12.

[16] BubalikM., BeckA., Baladincz J., HancsokJ., **(2009)**, Development of deposit control additives for diesel fuel, *Petroleum & Coal*, **51(3)**, 167-175

[17] ShuklaS. K., Tiwari V.K., Rani S., Ravikan, Tewari I.C., **(2011)**, Studies on synthetic, biological and insecticidal aspectsof somediorganotin (IV) diamides, *International Journal of Agriculture Sciences*, **3(3)**, 121-127.

[18] Butler D. N., Shang M., David M., Johnston M. R., **(2001),** Aryl ring dynamics in bis-succinimido- cyclophanes,*ARKIVOC*, **(xii)**, 27-34.

[19] Basu P. K., GhoshA., **(2012)**,Regioselective 5-exo-Trig Heterocyclization of2-Allyl-1-naphthols under the Influence ofN-Iodosuccinimide orMolecular Iodine inAqueousMicelle, *Hindawi Publishing Corporation Organic Chemistry International*, *Volume 2012*, **(Article ID 810476)**, 1-7

[20] Ali Z. Y., Mohamed S. A., EdreesM. M., **(2012),**Biological Screening for Some Newly Synthesized Pyrazolo[3,4-d]Pyrimidine Derivatives, *Nature and Science*, **10(7),** 1-12.

[21] WangaY. S., ChangaN. C.,LinbC. H., TaibH. M.**,(2008)**, Efficient Synthesis of 4,5-Disubstituted-3-toluenesulfonyl Glutarimides. Application to the Formal Synthesis of Mappicine Ketone, *Journal of the Chinese Chemical Society*,**(55)**, 431-434.

[22] Benjamin E., HijjiY., **(2008**), The Synthesis of Unsubstituted Cyclic Imides Using Hydroxylamine under Microwave Irradiation,*Molecules*, **(13)**, 157-169.

[23] ShetgiriN.P., NayakB.K., **(2005)**, Synthesis and antimicrobial activity of some succinimides, *Indian Journal of Chemistry*, **44B**, 1933-1936.

[24] Vitnik V. D., Popovik-Dordevikc J. B., Vitnik Z. J., Ivanoic M. D., Juranic I. O., **(2012),** Rotational Isomers, NMR and Reactivity Analysis of tert- pentyl-1-benzyl-4-methyl-2,6-glutarimide-3-carboxylate(PBMG), *University of Belgrade, Serbaria*, **(B-06-P)**, 88-90.

[25] Fortuna C. G., Barresi V., MussoN., MusumarraG., **(2009)**,Synthesis and applications of new trans1-indolyl-2-(1-methyl pyridinium and quinolinium-2-yl)ethylenes,*ARKIVOC*, **(viii)**, 222-229.

[26] Blanco M. M., Salerno A., PerilloI. A., **(2010)**, Mass spectrometry ofaromatic cyclic imides and amides. Part II: electron ionization induced decomposition of N-substituted 3,4-pyridinedicarboximides,*ARKIVOC*, **(xi)**, 215-231.

[27] Alaa A. M., Abdel A., **(2007)**, Novel and versatile methodology for synthesis of cyclic imides andevaluation of theircytotoxic, DNA binding, apoptotic inducingactivities and molecular modeling study,*European Journal of Medicinal Chemistry*, **(42)**, 614-626.

[28] Kossokowski J., Javocka-WierzbaM., **(2003)**, Synthesis of new N-substituted cyclic imides with an expected anxiolytic activity.XII.derivatives of 1-methoxy-5-bicyclo [2.2.2]-oct-5-one-2,3-dicarboximide, *Acta Poloniae Pharmaceutical Drug Research*, **60(5)**, 367-374.

[29] TaherpourA. A., Kheradmand K., **(2008)**, One-PotMicrowave Assisted Solid Phase Synthesis of Cyclic Imides from Cyclic Anhydrides,*Asian Journal of Chemistry*,**20(5)**, 3341-3344.

[30] KodeN. R., Jean J. V. E.,Mayence A., Wang G.,Huang T. L., **(2013)**, Design and Synthesis of N1,N5-bis[4-(5-Alkyl-1,2,4-oxadiazol-3-yl)phenyl]glutaramides as Potential Antifungal Prodrugs,*Molecules*, **(18)**, 11250-11263.

[31] UlianasA., Heng L. Y., Ahmad M., **(2012)**,A Biosensor for Urea fromSuccinimide-Modified AcrylicMicrospheres Based on Reflectance Transduction, *Sensors*,**(11)**, 8323-8338.

[32] Prezhdo V.V., Kowalski P., Kowalska T., Zubkova V.V., Olan K., Prezhdo O.V., **(2011)**,Molecular polarizability anisotropy of some five-membered cyclic imides, *Journal of Molecular Structure*, **(997)**, 20-29.

[33] Rajput A.P., Girase P.D., **(2012)**, Synthesis of 2,6-diazido-1-(N-substitutedphenyl)-1,4-dihydropyridine- 3,5-dicarbaldehydes and their transformationinto multivariant functionalities, *International Journal of ChemTech Research*, **4(2)**, 634-639.

[34] SharmaR.N., Sharma K.P., DixitS.N., **(2010)**, Synthesis, Characterization, and Biological activities of Some New Arylazopyrazoles, *International Journal of Chem Tech Research*, **2(2)**, 800-806.

[35] Rajput A. P., Rajput S. S., **(2009)**, Synthesis and Microbial Screening of SevenMemberedHeterocyclic ring Compounds from 1, 2-Diaminobenzene, International *Journal of PharmTech Research*, **1(3)**, 900-904.

[36] RajputS.S., **(2012)**, Synthesis and Characterization of Bis-Heteroyclic Derivatives of 1-(3-Chlorophenyl)-Pyrrolidine-2, 5-Dione, *International Journal of Advances in Pharmacy, Biology and Chemistry*, **1(2)**, 242-246.